

GRB 060117: Reverse + forward shock solution

MARTIN JELÍNEK⁽¹⁾ (*), MICHAEL PROUZA⁽²⁾⁽³⁾⁽⁵⁾, PETR KUBÁNEK⁽⁴⁾,
RENÉ HUDEC⁽⁴⁾, MARTIN NEKOLA⁽⁴⁾, JAN ŘÍDKÝ⁽³⁾⁽⁵⁾ and JIŘÍ GRYGAR⁽³⁾⁽⁵⁾

⁽¹⁾ *Instituto de Astrofísica de Andalucía, 180 08 Granada, Spain*

⁽²⁾ *Nevis Institute and Department of Physics, Columbia University, New York, NY 10027, USA*

⁽³⁾ *Fyzikální ústav AV ČR, Na Slovance 2, Praha 8,*

⁽⁴⁾ *Astronomický ústav AV ČR, Fričova 298, Ondřejov, Czech Republic*

⁽⁵⁾ *for the Pierre Auger Collaboration*

Summary. — We present a discovery and observation of an extraordinarily bright prompt optical emission of the GRB 060117 obtained by a wide-field camera atop the robotic telescope FRAM of the Pierre Auger Observatory from 2 to 10 minutes after the GRB. We found rapid average temporal flux decay of $\alpha = -1.7 \pm 0.1$ and a peak brightness $R = 10.1$ mag. We interpret the shape of the lightcurve as a transition between reverse and forward shock emission.

PACS 95.55.Cs – Ground based ultraviolet, optical and infrared telescopes.

PACS 95.75.De – Photography and photometry.

PACS 98.70.Rs – Gamma-ray sources; gamma-ray bursts.

1. – Swift detection

A bright long-soft GRB 060117 was detected by *Swift* satellite on January 17, 2006, at 6:50:01.6 UT. It showed a multi-peak structure with $T_{90} = 16 \pm 1$ s with maximum peak flux 48.9 ± 1.6 ph cm⁻²s⁻¹. Thus, GRB 060117 was — in terms of peak flux — the most intense GRB detected so far by Swift Coordinates computed by *Swift* were available within 19 s and immediately distributed by GCN [1].

2. – FRAM and optical transient observation

FRAM is part of the Pierre Auger cosmic-ray observatory [4], and its main purpose is to immediately monitor the atmospheric transmission. FRAM works as an independent,

(*) mates@iaa.es




Fig. 1. – The R-band afterglow lightcurve of GRB 060117. The lightcurve is fitted as a superposition of reverse shock (dotted line) and forward shock (dashed line).

RTS2-driven, fully robotic system, and it performs a photometric calibration of the sky on various UV-to-optical wavelengths using a 0.2m telescope and a photoelectric photomultiplier.

FRAM received the notice at 06:50:20.8 UT, 19.2s after the trigger and immediately started the slew. The first exposure started at 06:52:05.4, 123.8s after the GRB. Eight images with different exposures were taken before the observation was terminated. A bright, rapidly decaying object was found, and its presence was reported by [3] soon after the discovery. The FRAM lightcurve for this optical transient is in Figure 1.

3. – Interpretation

Our preferred interpretation (based on the work of [5]) is to fit the data as a transition between the reverse and the forward shock with the passage of the typical frequency break ν_m through the observed passband at time $t_{m,f}$. Corresponding decay indices are $\alpha_{Reverse}=2.49\pm0.05$ and $\alpha_{Forward}=1.47\pm0.03$ (see Fig 3).

Other possible interpretations and more details about FRAM telescope, data processing and other follow-up attempts can be found in [2].

* * *

The telescope FRAM was built and is operated under the support of the Czech Ministry of Education, Youth, and Sports through its grant programs LA134 and LC527. MJ would like to thank to the Spanish Ministry of Education and Science for the support via grants AP2003-1407, ESP2002-04124-C03-01, and AYA2004-01515 (+ FEDER funds), MP was supported by the Grant Agency of the Academy of Sciences of the Czech Republic grant B300100502.

REFERENCES

- [1] CUMMINGS, J., BARBIER, L., BARTHELMY, S., ET AL., GCN 4538, 2006
- [2] JELÍNEK, M., PROUZA M., KUBÁNEK, P. ET AL., Astron. Astrophys. 410, L119, 2006
- [3] KUBÁNEK, P., JELÍNEK, M., PROUZA, M., NEKOLA, M., and HUDEC, R., GCN 4535, 2006

- [4] PIERRE AUGER COLLABORATION, Proceedings of the 29th International Cosmic Ray Conference, Pune, India, also the FERMILAB preprint FERMILAB-PUB-05-469-A-TD, 2005
- [5] SHAO, L. & DAI, Z. G., *Astrophys. J.* 633, p. 1027, 2005